Volgograd state medical university



# Department of pathological anatomy

#### Lecture

## INFLAMMATION

Dentistry faculty

Inflammation is a protective response intended to eliminate the initial cause of cell injury as well as the necrotic cells and tissues resulting from the original insult

## **Etiology of Inflammation**

## **Exogenous factors**

#### **Physical agents:**

- extreme temperatures,
- electricity,
- irradiation,
- mechanical injures, etc.

#### **Chemical agents:**

- acids,
- alkalis,
- drugs.

#### **Biological agents:**

- Microorganisms (bacteria, viruses, fungi),
- parasites (helmints)
- Insects/

#### **Endogenous factors**

- immune cells and complexes
- tissue necrosis
- products of metabolism,

# Inflammatory phases

1. Alteration – damage\injury (dystrophy and necrosis) and production of mediators.

2. Exudation – the reaction of microcirculation, formation of liquid exudate, migration of leukocytes, phagocytosis.

 Proliferation - proliferation of cell of hematogenous (macrophages, lymphocytes) and histiogenous (fibroblasts) cells.

# Chemical mediators of inflammation

## Chemical mediators

- Plasma-derived:
  - Complement, kinins, coagulation factors
  - Many in "pro-form" requiring activation (enzymatic cleavage)
- Cell-derived:
  - Preformed, sequestered and released (mast cell histamine)
  - Synthesized as needed (prostaglandin)


## Inflammatory mediators

# Plasma-derived (circulating) mediators:

- 1) Kallikrein-kinin system:
- Kallikrein,
- Kinins (bradykinin);
- 2) The complement system (C5a, C3a, C3b, complex C5a C9a, C5a des Arg);
- 3) Blood coagulation system (Hageman factor),
- 4) Fibrinolytic blood system.

#### Cell-derived (local) mediators:

- 1) Vasoactive amines:
- Histamine,
- Serotonin,
- 2) Arachidonic acid metabolites:
- Prostaglandins,
- Leukotrienes,
- 3) PAF (Platelet Activation Factor),
- 4) *Cytokines:* IL-1, TNF (a, b), IL-8, g-interferon;
- 5) Contents of lysosomes.



© Elsevier. Kumar et al: Robbins Basic Pathology 8e - www.studentconsult.com


#### **Diversity of Effects of Chemical Mediators**

#### **Prostaglandins:**

- Vasodilation
- · Pain
- Fever
- Potentiating edema


#### IL-1 and TNF:


- Endothelial-leukocyte interactions
- Leukocyte recruitment
- Production of acute-phase reactants



### **Basophils & Mast Cells**

#### Histamine





### Role of Mediators in Different Reactions of Inflammation

Vasodilatation	Prostaglandins Nitric oxide Histamine	
Increased vascular permeability	Vasoactive amines C3a and C5a (through liberating amines) Bradykinin Leukotrienes C4, D4, E4 PAF Substance P	
Chemotaxis, leukocyte recruitment and activation	C5a Leukotriene B4 Chemokines IL-1, TNF Bacterial products	
Fever	IL-1, TNF Prostaglandins	
Pain	Prostaglandins Bradykinin	
Tissue damage	Neutrophil and macrophage lysosomal enzymes Oxygen metabolites Nitric oxide	

## **Summary of Mediators of Acute Inflammation**

	ACTION				
Mediator	Source	Vascular Leakage	Ch	Chemotaxis Other	
Histamine and serotonin	Mast cells, platelets	+	85.54		
Bradykinin	Plasma substrate	+	-	Pain	
C3a	Plasma protein via liver	1.4	-	Opsonic fragment (C3b)	
C5a	Macrophages	+	+	Leukocyte adhesion, activation	
Prostaglandins	Mast cells, from membrane phospholipids	Potentiate other mediators	•	Vasodilatation, pain, fever	
Leukotriene B <sub>4</sub>	Leukocytes	•	+	Leukocyte adhesion, activation	
Leukotrienes C <sub>4</sub> D <sub>4</sub> E <sub>4</sub>	Leukocytes, mast cells	1. <b>4</b> 1	(i=:)	Bronchoconstriction, vasoconstriction	
Platelet Activating Factor (PAF)	Leukocytes, mast cells	+	+	Bronchoconstriction, leukocyte priming	
IL-1 and TNF	Macrophages, other	30 <b>%</b> 3	+	Acute-phase reactions, endothelial activation	
Chemokines	Leukocytes, others		+	Leukocyte activation	
	Macrophages, endothelium	.+	+	Vasodilatation, cytotoxicity	

Acute inflammation has two major components

- Vascular changes
- Cellular events

## Vascular

- Changes in Vascular Caliber and Flow
- Increased Vascular Permeability

# Pathogenesis of exudation phase

- 1. Initial short-term microvascular spasm, followed be a long persistent dilation
- 2. Release of plasma and its protein components (albumin, globulins, fibrinogen)
- 3. Stasis, diapedesis of leukocytes and red blood cells in the surrounding tissue
- 4. Formation of exudate
- 5. Phagocytosis



lasting few seconds


arteriolar vasodilation




increased viscosity & slowing of circulation








## Increased Vascular Permeability



## **Cellular Events**


 an important function of the inflammatory response is to deliver leukocytes to the site of injury and to activate them

### Эмиграция нейтрофилов




# EXTRAVASATION of PMNs

- MARGINATION (PMN's go toward wall)
- ROLLING (tumbling and HEAPING)
- ADHESION
- TRANSMIGRATION (DIAPEDESIS)



## Leukocyte Recruitment

- Margination: Slow blood flow
  - White cells redistribute along periphery of lumen



## Leukocyte recruitment

#### **MARGINATION**





## Chemotaxis

- Leukocytes follow chemical gradient to site of injury (chemotaxis)
  - Soluble bacterial products
  - Complement components (C5a)
  - Cytokines (chemokine family e.g., IL-8)
  - LTB<sub>4</sub> (AA metabolite)( lymphotoxin beta -LTB,AA-Arachidonic acid)
- Chemotactic agents bind surface receptors inducing calcium mobilization and assembly of cytoskeletal contractile elements

# Types of exudate

- 1. Serous exudate.
- 2. Fibrinous exudate.
- 3. Purulent (suppurative) exudate.
- 4. Hemorrhagic exudate.
- 5. Mixed exudate.


## **PHAGOCYTOSIS**


- The process of engulfment of solid particulate material by the cells.
- · 2 main types of phagocytic cells
  - Polymorphonuclear neutrophils (PMNs): early in acute inflammatory response, also known as microphages
  - Macrophages: Circulating monocytes and fixed tissue mononuclear phagocytes
- This phagocytic cells releases proteolytic enzymes
   lysozyme, protease, collagenase, elastase,
  - lipase, proteinase, gelatinase and acid hydrolases


# **Phagocytosis**


Consists of 3 distinct but interrelated steps

- recognition and attachment of the particle to the ingesting leukocyte
- engulfment, with subsequent formation of a phagocytic vacuole
- killing and degradation of the ingested material.









## Purpose of proliferative phase

 Completing of phagocytosis - final destruction and elimination of the etiologic agent

2. Delimitation of inflammatory focus from healthy tissue

Complete regeneration/Replacement lesion scar tissue

# Types of inflammation

- Acute inflammation lasts from several days up to several months – in the focus of inflammation - neutrophils, intravascular platelet activation – Exudative inflammation and rarely observed productive (viruses)
- Subacute inflammation lasts from several weeks up to several months – in the focus of inflammation - neutrophils, lymphocytes, plasmocytes, macrophages (approximately in equal proportions) – exudative-productive inflammation
- Chronic inflammation lasts from a few months up to tens of years – alternating exacerbations and remissions – in the focus of inflammation – mononuclear cells (lymphocytes, plasmocytes, macrophages), in case of exacerbations neutrophils are added – productive inflammation, during exacerbations an exudative reaction is added – presence of fibrosis is essencial

## SIGNS OF INFLAMMATION

- RUBOR- REDNESS DUE TO INCREASED BLOOD BLOW AND VASODILATION
- CALOR- OR HEAT DUE TO INCREASE BLOOD FLOW TO THE PERIPHERY
- TUMOR- SWELLING FROM INFLAMMATORY EDEMA
- DOLOR-PAIN FROM SWELLING AND PRESENCE OF INFLAMMATORY MEDIATORS
- FUNCTIO LAESA-LOSS OF FUNCTION DUE TO MAIN

AND STRUCTURAL NECROSIS

# Systemic effects

- Fever
  - One of the easily recognized cytokine-mediated (esp. IL-1, IL-6, TNF) acute-phase reactions including
    - Anorexia
    - Skeletal muscle protein degradation
    - Hypotension
- Leukocytosis
  - Elevated white blood cell count

	TRANSUDATE	EXUDATE	
COLOR	Clear, water-like, or pale yellow of plasma	Cloudy, white, yellow or red	
CONSISTENCY	Thin and watery, no tissue fragments	Thick and creamy, con- tains tissue fragments	
ODOR	None	May have an odor	
рН	Alkaline	Acid	
SPECIFIC GRAVITY	1.015 or lower	1.018 or higher	
FROTEIN CONTENT	Low, less than 3%	High, more than 4%	
CELL COUNT	Low; none or few WBC and RBC	High; many WBC and RBC	
ENZYME CONTENT	Low	High	
BACTERIA	None	May be present	
INFLAMMATION	None present	Associated with Inflammation	

# Types of inflammation

Exudative infl.

 Proliferative (productive) infl.

## Types of acute inflammation

- 1. Serous inflammation;
- 2. Fibrinous inflammation → croupous;diphtheritic;
- 3. Purulent inflammation abscess, phlegmon, empyema, furuncle, carbunckle;
- 4. Hemorrhagic inflammation
- 5. Catarrhal inflammation
- 6. Mixed inflammation

# MORPHOLOGICAL PATTERNS OF ACUTE INFLAMMATION

- SEROUS INFLAMMATION
- Outpouring of thin fluid
- Depends on secretion of mesothelial cells peritoneal pleural, pericardial cavities- effusions
- Skin blister- viral ,burns- large accumulation of serous fluid
- Watery, protein-poor effusion (e.g., blister)
- FIBRINOUS INFLAMMATION
  - Fibrin accumulation
  - Either entirely removed or becomes fibrotic
- Inflammation in the lining of body cavities meninges pericardium, pleura
- Scarring

#### Morphologic Patterns

#### Serous Inflammation

Inflammation of serous membrane characterized by clear fluid in serous cavity (pleural, peritoneal pericardial & synovial cavities)


E.g. skin Blisters caused by Burns OR viral infection

## Morphologic Patterns




CONTACT DERMATITIS BLISTERS FROM THE POISON IVY PLANT'S TOXIC PRINCIPLE, URUSHIOL. THIS IS A CLASSIC EXAMPLE OF A SEROUS EXUDATE.

#### Serous inflammation



#### Herpes simplex:

 Vesicles of the skin;



 Multi-nuclear cells and intranuclear inclusions.


#### Serous inflammation





Chickenpox



#### Serous inflammation







#### Morphologic Patterns

#### Fibrinous Inflammation


- Severe injury with excessive deposition of Fibrin in serous cavity
- Exudate and Fluid is removed by lymphatic
- Fibrinous exudate may be degraded by Fibrinolysis and removed by macrophage resulting in Resolution.
- Incomplete Removal of fibrin resulting in organization and scarring with
- Fibrous Adhesion of pleura OR pericardium.

#### **Fibrinous Inflammation**

1. Fibrinous Parenchymal Inflammation

2. Fibrinous Serosal Inflammation

3. Fibrinous Mucosal Inflammation



#### Fibrinous inflammation

#### **Croupous Type**


- an easily removable pseudomembrane covering the necrosis, which is limited to the mucosal epithelium
- Expl: Diphtheric laryngotracheitis

#### **Diphtheritic Type**

- necrosis extending into the submucosa,
- a wide area of fibrinous exudate is covered by an adhesive pseudomembrane that can only be forcibly removed.
- Expl: Diphtheric tonsillitis and pharyngitis, • Dysentery,
  • Antibiotic associated colitis (pseudomembranous colitis)




# Fibrinous pericarditis



# Fibrinous pericarditis

#### FIBRINOUS INFLAMMATION-PERICARDIUM



## Morphologic Patterns

#### Pseudomembraneous Inflammation

- Very severe ulcerative inflammation of mucous membranes
- Extensive Necrosis of surface epithelium
- Severe acute Inflammation of underlying tissue
- Pseudo membrane, consisting of exudate, fibrin, neutrophils RBC, Bacteria and tissue debris
- e.g. Diphtheria Larynx . pseudomembraneous Colitis clostridium difficile

## Morphologic Patterns


Pseudomembraneous Inflammation



### **Fibrinous inflammation**



#### **Fibrinous inflammation**




#### Outcomes of fibrinous inflammation

- 1. Resolution of exudate.
- Formation of adhesions in body cavities up to obliteration of cavities.

#### The value of fibrinous inflammation:

- This is very large, since it constitutes the morphological basis of diphtheria, dysentery, is observed with intoxications (uremia).
- With the formation of films in the larynx, the trachea the danger of asphyxia; when tearing films in the intestine bleeding from the resulting ulcers.
- Adhesive pericarditis and pleurisy are accompanied by the development of pulmonary heart failure.

#### Outcomes of fibrinous inflammation



 Formation of adhesions in body cavities up to obliteration of cavities

#### **Purulent Inflammation**

 General definition: Inflammation with exudate consisting primarily of died neutrophils and cellular debris (detritus).

# Types of purulent (suppurative) inflammation

- 1. Abscess (acute and chronic);
- 2. Phlegmon (hard and soft);
- 3. Empyema;
- 4. Furuncle;
- 5. Carbuncle.

#### **Abscess**

 local purulent inflammation with the formation of a cavity filled with pus;


# The wall of acute abscess consists of 2 layers:


- 1) the inner layer (granulation tissue) is a pyogenic membrane;
- 2) the outer layer compressed surrounding healthy tissue;

# The wall of the chronic abscess consists of 3 layers:

- 1) inner layer purulent-necrotic masses;
- 2) the middle layer is a granulation tissue;
- 3) outer layer connective tissue capsule.

## **ABSCESS-HEART**






# **Purulent** inflammation

Abscesses of the lung;

#### Abscess of the brain








Metastatic kidney abscesses



kidney abscesses




#### Periodontal Abscess




### Paratonzillar abscess



#### **Purulent inflammation**



 Abscess of the lung;




Abscessed pneumonia.

#### **Empyema**

- Definition: Suppurative inflammation in a body cavity.
- Pathogenesis: An empyema usually occurs when a suppurative inflammation of an organ breaks through into an adjacent cavity.
- Examples:
- Pericardial, peritoneal, and pleural empyema
- Gallbladder and appendiceal empyema;
- Middle ear and nasal sinus empyema;
- Pyosalpinx (pus in the uterine tube);
- Pyocephalus (pus in the cranial cavity);
- Hypopyon (pus in the anterior chamber of the eye).

## Purulent peritonitis





# Purulent pericarditis

## Empyema of the gallbladder





#### **Phlegmon**

 Definition: Diffuse suppurative inflammation that spreads primarily in loose fibrous connective tissue without sharp demarcation.

#### Examples:

- Muscular phlegmon;
- Phlegmon of the floor of the mouth;
- Mediastinal phlegmon;
- Phlegmon of the walls of hollow organs (such as phlegmon in cholecystitis, appendicitis).
- Erysipelas, inflammation in the connective tissue of the skin usually caused by beta-hemolytic streptococci involving a map-like pattern of erythema and swelling;

#### **Phlegmon**

- Diffuse (diffuse) purulent inflammation;
- It develops in the subcutaneous fatty cellulose, cellulose surrounding the internal organs;
- Solid phlegmon is characterized by the presence of foci of coagulation necrosis, which do not undergo melting, and are gradually rejected.
- Soft phlegmon is characterized by a lack of visible foci of necrosis in the tissues;


#### **Phlegmon**


- Paronychia is an acute purulent inflammation of the periosseous tissue.
- Panaritium acute purulent inflammation of the subcutaneous tissue of the finger.
- Phlegmon neck acute purulent inflammation of the cellulose neck, develops as a complication of pyogenic infections of the tonsils, maxillofacial system.
- Danger of phlegmon neck a purulent process can spread to the mediastinal fiber (purulent mediastinitis), pericardium (purulent pericarditis), pleura (purulent pleurisy).
- Phlegmon is always accompanied by severe intoxication and may be complicated by sepsis.

- Mediastinitis is an acute purulent inflammation of the mediastinal fiber.
- Distinguish: anterior and posterior purulent mediastinitis.
- Anterior mediastinitis is a complication of suppurative inflammatory processes of the anterior mediastinum organs, pleura, phlegmon of the neck.
- Posterior mediastinitis is most often caused by the pathology of the esophagus: for example, traumatic injuries caused by foreign bodies (the bone bone is especially dangerous), decaying esophagus cancer, purulent necrotic esophagitis, etc.
- Purulent mediastinitis is a very severe form of purulent inflammation, accompanied by severe intoxication, which is the cause of the patient's death.

- Paranephritis is a purulent inflammation of the perineal tissue.
- is a complication of purulent nephritis, septic infarction of the kidneys, decaying kidney tumors.
- Meaning: intoxication, peritonitis, sepsis.
- Parametritis is a purulent inflammation of the periarticular tissue.
- It occurs in septic abortions, infected labor, decay of malignant tumors.
- First purulent endometritis, then parametrite.
- Meaning: peritonitis, sepsis.
- Paraproctitis is the inflammation of the tissue surrounding the rectum.
- Causes: dysentery ulcers, ulcerative colitis, decaying tumors, anal fissures, hemorrhoids.
- Meaning: intoxication, the appearance of pararectal fistulas, the development of peritonitis.








# Phlegmonous appendicitis



# Phlegmonous appendicitis




### **Purulent mediastinitis**



# Odontogenic phlegmon of face and neck





### **Furuncle**

- this is an acute purulent-necrotic inflammation of the hair follicle (follicle) and associated with it the sebaceous gland with surrounding fiber.
- Causes: Staphylococcus aureus, streptococcus.
- Conditions conducive to the development of furuncle:
- 1. constant contamination of the skin and rubbing clothes,
- irritation with chemicals, abrasions, combs and other microtraumas,
- 3. intensified activity of sweat and sebaceous glands, beriberi,
- 4. metabolic disorders (eg, diabetes mellitus),
- 5. starvation,
- 6. weakening of the body's defenses.



# **Furuncle**





# ıncle



### Carbuncle

- this is an acute purulent inflammation of several nearby arranged hair follicles and sebaceous glands with necrosis of the skin and subcutaneous tissue of the affected area.
- Occurs: when pyogenic microbes enter the ducts of the sebaceous or sweat glands, as well as when they penetrate the skin through minor lesions, squeezing out the furuncle.
- The conditions of development and localization are the same as in the furuncle.
- Macroscopically: an extensive dense, red-crimson infiltration on the skin, in the center of which there are several purulent "heads".

## Carbuncle

- The most dangerous carbuncle of the nose and especially the lips, in which a purulent process can spread to the membranes of the brain, resulting in purulent meningitis.
- Treatment operative; at the first symptoms of the disease you need to consult a surgeon.
- Value. Carbuncle is more dangerous than a boil, is always accompanied by severe intoxication.
- Complications: purulent lymphadenitis, purulent thrombophlebitis, erysipelas, phlegmon, sepsis.

# Carbuncle









## Outcomes of purulent inflammation

- Formation of external (in the external environment) and internal (in some kind of hollow organ) fistula;
- "Purulent fouling" (blind pockets with pus, departing from the focus of purulent inflammation);
- 3. Formation of sequestration (sequestration);
- 4. Scar formation (organization) after evacuation of the purulent inflammation site;
- 5. Formation of the capsule (encapsulation).

## **Hemorrhagic Inflammation**


- Definition: Exudative inflammation involving microvascular injury with massive microvascular bleeding, producing an exudate with a high erythrocyte content.
- Biologic purpose: Exudative inflammation due to severe vascular injury.
- Morphology: The inflamed area is usually necrotic and filled with blood.
- Etiologic factors:
- bacterial exotoxins and endotoxins;
- viral cytopathic effect on endothelium;
- proteolytic tissue destruction;
- cytotoxic injury in hypersensitivity type III.

# **Hemorrhagic Inflammation**

### **Examples:**


- Plague
- Influenzal Pneumonia
- Disorders Associated with Enterohemorrhagic E. Coli
- Anthrax
- Viral Hemorrhagic Fever
- Acute Pancreatitis

### **HEMMORRHAGIC INFLAMMATION-RIGHT**






# Hemorrhagic Inflammation



# Hemorrhagic pancreatitis








 Hemorrhagic form of herpes

### **Catarrhal Inflammation**


- Definition: Exudative inflammation occurring exclusively on the mucous membranes of the respiratory and gastrointestinal tracts and producing a watery exudate of serum and mucus.
- Subtypes: mucus (most frequent), serous, purulent, hemorrhagic
- Etiologic factors:
- hypersensitivity reactions;
- bacterial and viral tissue injury;
- physical and chemical tissue injury.


#### Catarrhal tracheitis






Catarrhal sinusitis









Catarrhal colitis



#### Catarrhal tracheitis





### **Mixed Inflammation**

- In those cases when one exudate is joined by another, mixed inflammation is observed.
- serous-fibrinou inflammations,
- purulent-hemorrhagic inflammations,
- fibrinous-hemorrhagic inflammation.
- Most often, a change in the type of exudative inflammation is observed with the addition of a new infection, a change in the reactivity of the organism.

## Outcomes of acute inflammation

#### **Favorable:**

- Absorption,
- tissue repair (the most favorable outcome)
- Organization scar formation

#### **Unfavorable:**

- Acute organ insufficiency
- Abscess formation pyonecrotic cavity
- Persistence of inflammation and chronicity

# Possible outcomes of acute inflammation

- Complete resolution
  - Little tissue damage
  - Capable of regeneration & restoration of injury cell to normal
  - Resolution involves neutralization, spontaneous decay of chemical mediators, subsequent return of normal vascular permeability, cessation of leukocyte infiltration, death by apoptosis removes edema, protein, foreign substance & necrotic debris.
- Scarring (fibrosis)
  - In tissues unable to regenerate
  - Excessive fibrin deposition organized into fibrous tissue
  - in many pyogenic infection intense neutrophil infiltration
     &liquefaction of tissue pus formation- fibrosis.

# Outcomes (cont'd)

- Abscess formation occurs with some bacterial or fungal infections
- Pneumonia, chronic lung abscess, peptic ulcer of duodenum or stomach –persist months or yrs
- Progression to chronic inflammation (next)

## **Chronic Inflammation**

 is inflammation of prolonged duration (weeks to months to years) in which active inflammation, tissue injury, and healing proceed simultaneously.

### Chronic Inflammation

- Inflammation for a prolonged period (weeks-months)
- Onset
  - After Acute Inflammation
    - When injurious agent is not removed promptly
    - More and more inflammatory cells are recruited to site of injury
  - Insideous Onset
    - Low intensity injury for long period of time
    - Inflammatory response is not overwhelming
    - Injury and inflammation persist

Chronic inflammation arises in the following settings:

- T lymphocyte-mediated immune response called delayed-type hypersensitivity
- 2. Immune-mediated inflammatory diseases
- 3. autoimmune diseases

### Chronic Inflammation - Causes

- Persistent infection
  - МТВ
  - Viruses
  - Fungi
  - Parasites
- Immune mediated
  - Autoimmune diseases (Rheumatoid arthritis)
  - Allergic diseases (Broncial asthma)
- Prolonged exposure
  - Silicosis (Pulmonary fibrosis)

### Chronic Inflammation - Features

- Inflammatory infiltrate
  - Macrophages
  - Lymphocytes
  - Plasma Cells
- Tissue destruction
  - Persistent injury
  - Inflammatory cells and Mediators
- Repair
  - Angiogenesis
  - Fibrosis

### Chronic Inflammation - Cells

- Macrophages
- Lymphocytes
- Plasma Cells

- Eosinophils
- Mast Cells
- Neutrophils!

## **Chronic inflammation**

# Chronic productive inflammation:

- 1. Interstitial inflammation;
- 2. Granulomatous inflammation;
- 3. Inflammation with the formation of polyps and genital warts.

# Chronic exudative inflammation:

- 1. Chronic abscess;
- 2. Chronic catarrhal inflammation.

## The main types of productive (proliferative) inflammation

1. Interstitial (interstitial) inflammation;

2. Granulomatous inflammation;


3. Inflammation with the formation of polyps and genital warts.

### Interstitial Inflammation

- Characteristic: the formation of focal or diffuse inflammatory cell infiltrate in the stroma of parenchymal organs myocardium, liver, kidneys, lungs.
- The main cells: lymphocytes, histiocytes, plasma cells, single neutrophils, eosinophils and mast cells.
- In the parenchymal elements dystrophic and necrobiotic changes.


- Examples: acute and chronic interstitial pneumonia, idiopathic myocarditis Abramov-Fiedler, chronic interstitial nephritis during chronic pyelonephritis.
- Outcome is sclerosis.

### Interstitial inflammation



Interstitial pneumonitis

### Interstitial myocarditis



## Inflammation with the formation of polyps and genital warts

- Productive inflammation, in which there is a simultaneous proliferation of parenchymal elements and stroma;
- The stroma is diffusely infiltrated by mononuclears.
- Characteristic of chronic course;
- Typical localization is the mucous membranes.
- Examples: polyps of the nasal cavity, maxillary sinuses, bronchi, gastric mucosa, intestine, uterus.

### Polyp of intestine




 Pointed condylomas - formed at the junction of the flat and prismatic epithelium (anus, genitalia);


are caused by papillomavirus (HPV);

 may be accompanied by dysplasia; are a risk factor for squamous cell carcinoma.


### **Genital Warts**



**Papillomavirus** 



Coilocyte reaction



Acanthosis, hyperkeratosis, parakeratosis, cytoplasmic vacuolation, koylocytosis

### **Granulomatous Inflammation**

- Distinctive type of chronic inflammation
- Attempt to contain an offending agent that is difficult to eradicate
- Strong activation of T cells and Macrophages
- Examples:
  - TB

- Fungal Infection

- Syphilis

- Leprosy

Sarcoidosis

- Crohn's

### GRANULOMATOUS INFLAMMATION

- Granulomatous inflammation occurs after the acute-phase response and consists of epithelioid and giant cells.
- Two types of granuloma:
  - Foreign Body granuloma-formed in response to indigestible materials
  - Allergic granuloma-formed in delayed hypersensitivity reactions

### **Granulomatous inflammation**

 Characterized by the predominance of activated macrophages and / or their derivatives;

Morphological substrate - granuloma;

 Granuloma is a focal cluster of cells capable of phagocytosis monocyte-macrophage nature.

### Etiology of granulomatous inflammation

### Endogenous factors:

poorly soluble products
 of damaged tissues,
 especially adipose
 tissue,

products of impaired metabolism (urate).

### Exogenous factors:

- biological (bacteria, fungi, protozoa, helminths);
- organic and inorganic substances (dust, fumes, medicines, etc.)

### Classification of granulomas (by etiology)

### Granulomas of known etiology

 Infectious granulomas (in typhus and typhoid fever, rabies, viral encephalitis, actinomycosis, schistosomiasis, tuberculosis, leprosy, syphilis, etc.)

# 2. Noninfectious granulomas (organic and nonorganic dusts - wool, flour, silicon oxide, asbestos, etc., foreign bodies, medicinal effects).


### Granulomas of unknown etiology

- 1. sarcoidosis,
- 2. Crohn's disease,
- primary biliary cirrhosis of the liver, etc.

### Morphogenesis of granuloma


- I stage. Accumulation in the lesion of tissue damage of young monocytic phagocytes.
- Il stage. Maturation of these monocytic phagocytes into macrophages and the formation of macrophagial granuloma.
- III stage. Maturation and transformation of monocytic phagocytes and macrophages into epithelioid cells and formation of epithelioid-cell granuloma.
- IV stage. Transformation of epithelioid cells into multinuclear giant cells and formation of giant cell granuloma.

#### **Granulomatous inflammation**




Epithelioid cells;


### Pirogov-Langhan's multinuclear giant cells



#### LANGHAN'S GIANT CELLS


### FOREIGN BODY GIANT CELL- IN SUTURE LINE






### Granuloma

- Focus of chronic inflammation
- May have central necrosis
- Aggregation of macrophages that are transformed into epithelioid cells
- Epithelioid cells fuse to form Giant Cells
- Surrounded by lymphocytes and plasma cells




### Foreign Body Granuloma

- Inert foreign bodies
  - Talc
  - Sutures




- Foreign Body-type Giant Cells
  - Haphazardly arranged nuclei


### Immune Granuloma - TB




### Immune Granuloma - Sarcoidosis



### SYNCYTHIAL GIANT CELLS



### GRANULOMATOUS INFLAMMATION WITH SPHERULES



### Classification of granulomas

### Immune granulomas

- Predominantly epithelioidcell;
- A large number of lymphocytes and plasma cells;
- in tuberculosis, leprosy, syphilis, sclera,
- in exposed to organic dust particles, aerosols containing the proteins of birds, fish, animal hair.

### Non-immune granulomas

- Primarily macrophagal or giant cell;
- around foreign bodies (suture material), animal parasites, particles of inorganic dust.

### Classification of granulomas (depending on the degree of rapidity of cell exchange within the granuloma)

### Rapidly updated granulomas

- Update for 1-2 weeks (cells quickly die and replaced with new ones);
- Granulomas produce very toxic substances (tuberculosis, leprosy);
- Predominantly epithelioidcell;
- Alien material is only partially located in macrophages.

#### Slowly updated granulomas

- The kinetics of cell metabolism is sharply delayed;
- Occur when exposed to inert, low-toxic substances;
- Predominantly giant-celled;
- The pathogen is entirely located in macrophages.

### **Outcomes of granulomas**

- Resolution of cellular infiltrate (a rare option, acute infections - rabies, abdominal and typhus);
- Sclerosis with the formation of a scar or fibrous nodule - most often;
- 3. Necrosis of granuloma tuberculosis granuloma, etc.
- 4. Suppuration of granuloma mycoses, sap, yersiniosis, tularemia.

### Tuberculosis granuloma

### Macroscopically:

- The diameter of the granulomas is 1-2 mm,
- Numerous confluent granulomas - small, millet-like tubercles -> miliary (from the Latin miliarius millet) tuberculosis.

### Microscopically:

- In the center a focus of caseous necrosis,
- Around a shaft of epithelioid cells,
- Further a shaft of lymphocytes with an admixture of giant multinucleated cells Pirogov-Langhans.
  - When impregnating with silver salts a network of argyrophilic fibers;
  - There are no blood vessels;
  - In giant cells when stained according to the Tsil-Nielsenumycobacterium tuberculosis.


### Granulomatous inflammation




 Miliary tuberculosis of the lungs / spleen



### **Granuloma in tuberculosis**



### **Granuloma in tuberculosis**



### Syphilitic granuloma (gumma)

 It is characteristic for the tertiary period of syphilis;

#### Macroscopically:

- Nodes with a diameter of 0.3-1.0 cm on the skin up to 4-5 cm in internal organs (bones, brain, liver, etc.);
- On the cut at the nodes jelly-like mass of yellow color (similar to glue gum arabic).


#### Microscopically:


- In the center there is a foci of caseous necrosis (larger than in tuberculosis);
  - In the periphery a lot of lymphocytes, plasma cells and fibroblasts;
  - characteristic the rapid formation of massive dense connective tissue (a kind of capsule);
- On the inside of the capsule are numerous small, and outside larger vessels with signs of productive endovascularity;
- Pale treponema is extremely rare (with Levadity silvering).


### Gumma in syphilis




### Gumma in syphilis










### **Syphilis**













Tertiary syphilis: Typical gummas. A. Wisdom. Color


Leprosy granuloma


• The causative agent is Mycobacterium Hansen-Neisser (coloration according to Tsil-Nielsen).


#### Cellular composition:


- 1. macrophages,
- 2. epithelioid cells,
- 3. giant cells,
- 4. plasma cells,
- Fibroblasts.
- Unfinished phagocytosis in the cytoplasm of macrophages -Mycobacterium Hansen-Neiser in the form of strictly ordered series (like cigarettes in a pack) -> increase in macrophages + appearance of fatty inclusions.
- Bonding of mycobacteria -> formation of leprosy balls
- Destruction of macrophages
- Phagocytosis of leprosy balls dropped by giant cells of foreign bodies.
- in 1 g of "blossoming" leprom 5x109 mycobacteria

### Leprosy granuloma










### Leprosy granuloma







### Granuloma in scleroma

Pathogen: wand of Volkovitch-Frish.

#### **Cellular composition:**

- Macrophages,
- 2. Lymphocytes,
- 3. Plasma cells,
- Eosinophilic corpuscles Roussel (products of degradation of plasma cells),
- Mikulich cells are specific cells very large singlenucleated macrophages with vacuolated cytoplasm + wolkovich-Frish sticks.


### Granuloma in scleroma


 Localization: the mucous membrane of the upper respiratory tract - nose, larynx, trachea, less often - bronchi.

 Outcome - sclerosis -> deformation of the mucosa - narrowing of the respiratory tract -> sometimes - asphyxiation.





# Scleroma





### **Granulomatous inflammation**







# Inflammation around foreign bodies:

 Aspiration pneumonia (inflammation around plant fibers) with giant cells of foreign bodies;

 Inflammation around the suture (stitches)

### **Granulomatous inflammation**





#### Granuloma in silicosis:

(macrophages, lymphocytes, plasmocytes, fibrosis);

#### Granuloma in sarcoidosis:

- "stamped" appearance,
- the central part from the epithelioid and giant multinucleated cells of Pirogov - Langhans,
- on the periphery lymphocytes, macrophages, plasma cells and fibroblasts.



## Actinomycosis

- systemic infection prone to sluggish, chronic course; characterized by the development of granulomas (actinomycetics), fistulas and abscesses.
- Pathogen various types of actinomycetes gram-positive bacteria of the genus Actinomyces, which can branch (form a mycelium).
- 3. Forms: cervico-facial, thoracic, abdominal and generalized.
- 4. Druses, surrounded by macrophages and neutrophils.





Figure 1.—Actinomycosis, jaw, observed at Letterman General Hospital, San Francisco, Calif., in a sergeant who had punctured the floor of his mouth with a weed stem while picking his teeth.




# Actinomycosis




### **Trichinosis**

- Helminthiasis from the group of nematodes,
- Characteristic: fever, myalgia, face swelling, skin rash, eosinophilia of the blood, and in severe cases - damage to internal organs and the central nervous system.
- Young larva through sarcolemma -> into muscle fiber, partially destroying it.
- Around the larva is a cellular infiltrate, and after 3-4 weeks.
   after the invasion a fibrous capsule with a network of blood vessels.
- The walls of the capsule gradually thicken, impregnated with calcium salts.
- The larvae remain viable for many years.




### **Trichinosis**






## **Trichinosis**








### • Насекомое Комар сосет кровь с личинками хоботном заносит личинок другой собане\* Личинки Сердечные ельминты Личинки http://eyorkie.ueoz.ru/pазитов


### **Dirofilariasis**

## **Dirofilariasis**






### **Dirofilariasis**



### Crohn's disease




 nonspecific disease of any part of the gastrointestinal tract.

 It is manifested by transmural inflammatory infiltration and sarcoid granulomas of the gastrointestinal tract.

## Crohn's disease








### Aspergillosis





Fungi - aspergillus.


 May complicate chronic diseases of the lungs such as bronchial asthma, chronic obstructive pulmonary disease.


### **Aspergillosis**




Ussually develops in patients with decreased immunological reactivity, caused by different diseases, genetic factors or due to treatment by cytostatics and immunosupressions.

#### Granuloma in aspergillosis









### Coccidiodomycosis

• H&e,





PAS-reaction.



















#### **Outcomes of chronic inflammation**

Fibrosis

Chirrosis

Chronic organ insufficiensy

#### Inflammation

#### Acute

- Rapid onset
- Short duration
- Neutrophils
- Exudation of fluid and plasma proteins
- Inflammation followed by Repair

#### Chronic

- Slow onset; may follow acute
- Longer duration
- Lymphocytes and Macrophages
- Proliferation of blood vessels
- Simultaneous Inflammation and Repair

